<http://www.diva-portal.org>

Postprint

This is the accepted version of a paper published in *Synlett: Accounts and Rapid Communications in Synthetic Organic Chemistry*. This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Citation for the original published paper (version of record):

Utas, J., Olofsson, B., Åkermark, B. (2006)

Efficient Synthesis of 2-Substituted Imidazoles by Palladium-Catalyzed Cross-Coupling with Benzylzinc Reagents.

Synlett: Accounts and Rapid Communications in Synthetic Organic Chemistry, (12): 1965-1967

<http://dx.doi.org/10.1055/s-2006-948162>

Access to the published version may require subscription.

N.B. When citing this work, cite the original published paper.

Permanent link to this version:

<http://urn.kb.se/resolve?urn=urn:nbn:se:su:diva-20420>

Efficient Synthesis of 2-Substituted Imidazoles by Palladium-Catalyzed Cross-Coupling with Benzylzinc Reagents

Josefin E. Utas, Berit Olofsson, Björn Åkermark*

Stockholm University, Department of Organic Chemistry, Arrhenius Laboratory, SE-106 91 Stockholm, Sweden.

Fax: +46-(0)8-154908; E-mail: Bjorn.Akermark@organ.su.se

Received:

Abstract: Substituted benzylzinc reagents have been used in novel cross-coupling reactions with 2-iodo imidazoles to form compounds containing both a phenol and an imidazole moiety. The intramolecular hydrogen bonding properties of these compounds were subsequently studied.

Key words: Cross-coupling, Palladium, Zinc, Benzylation, Heterocycles

Hydrogen bonding has recently been recognized to modify the redox properties and electron transfer rates of the amino acid tyrosine in certain enzymes.² This is of great importance in Photosystem II (PSII) in the photosynthesis, where a tyrosine residue (Tyr_Z) acts as an electron transfer mediator between the water-oxidizing manganese complex and the special pair of chlorophyll molecules called P680.¹ A histidine residue (His190) is also involved in this process, but the exact actions around the site remain unclear.³

As a model system for the interactions between Tyr_Z and His190 in PSII, we are currently investigating phenols where imidazole can act as a hydrogen bond acceptor and thus lower the phenol oxidation potential. According to preliminary density functional calculations,⁴ a strong intramolecular hydrogen bond could be expected in 7-membered rings such as compounds **1** (Figure 1). We thus designed structures **1a-c**, having various degrees of sterical hindrance, as target molecules for this study, and herein describe the synthesis and preliminary hydrogen bond measurements of these interesting compounds.

Figure 1 General structure of target compounds **1**.

As structures **1** contain a methylene bridge between the two aromatic rings, their synthesis is not trivial. Previous routes to similar compounds are sensitive to steric hindrance⁵ or involve several steps with limited possibilities to vary the substituents.⁶

Three different coupling strategies were thus investigated with test substrates to yield 2-benzyl-1-methylimidazole (**2**⁶, Scheme 1). Our first attempt involved the coupling of benzaldehyde with lithiated imi-

dazole **3**,^{7,8} but this strategy was hampered by problems in the subsequent reduction of the formed alcohol. Next, benzyl bromide was reacted with lithiated imidazole **3**, which resulted in a complicated mixture of products both in the presence⁹ and absence⁸ of PdCl₂(PPh₃)₃ (Scheme 1a).

To avoid the use of a basic lithium compound, which could deprotonate the relatively acidic benzyl halide,⁸ we looked into the possibility of employing a zinc reagent as transmetallating agent in a Negishi coupling.¹⁰ Benzylzinc reagents can be prepared by treating the corresponding benzyl halide with activated Zn dust,¹¹ and have successfully been employed in couplings with various halogenated heterocycles.¹²

Scheme 1 Strategies investigated towards imidazole **2**.

Benzyl bromide was thus converted to benzylzinc bromide (**4**¹³) and reacted with 2-iodo-imidazole **5a**¹⁴ in the presence of PdCl₂(PPh₃)₂ (Scheme 1b). Gratefully, the coupling could be smoothly accomplished at room temperature. An excess of zinc reagent **4** was used to assure full conversion, and compound **2** could be obtained in 68% yield.

Having found suitable coupling conditions, the benzylzinc reagents needed to form target compounds **1** were synthesized (Scheme 2). Starting from commercially available alcohol **6a**, the unsubstituted phenol derivative **7a**¹⁵ could be obtained in two steps, and *tert*-butyl derivative **7b** was formed from aldehyde **8** in a similar manner. The zinc reagents could conveniently be stored in the fridge until use.

Scheme 2. Synthesis of zinc reagents **7**.

Fortunately, the coupling of zinc reagent **7a** with 2-iodoimidazole **5a** in the presence of $\text{PdCl}_2(\text{PPh}_3)_3$ was successful, yielding compound **9a** in 62%. (Scheme 3). In an attempt to improve the yield, two other metal catalysts were employed in the coupling.¹⁰ $\text{Pd}(\text{OAc})_2$ together with tri(*o*-tolyl)phosphine resulted in 59% yield of **9a** whereas $\text{NiCl}_2(\text{dppp})_2$ only gave traces of the desired product. As the two Pd catalysts gave similar results, $\text{Pd}(\text{OAc})_2$ was chosen for further reactions due to easier purification.

Scheme 3. Synthesis of target molecules **1**.

To our delight, the $\text{Pd}(\text{OAc})_2$ -catalyzed coupling of *tert*-butyl-substituted zinc reagent **7b** with imidazole **5a** resulted in compound **9b** in excellent yield. Unfortunately, the coupling of **7a** with 1-benzylsubstituted imidazole **5b**⁸ was sluggish, and product **9c** could only be isolated in 25% yield together with recovered starting material.

Demethylation of the coupling products **9a,b** was accomplished with BBr_3 to yield the desired target molecules **1a** and **1b**.¹⁶

Finally, NMR studies were performed to elucidate whether hydrogen bonds were present in compounds **1a,b**. In CHCl_3 , the phenolic proton of 2,4-di(*tert*-butyl)phenol has a chemical shift of 4.62 ppm. Upon addition of 1-methylimidazole (1:1), *intermolecular* hydrogen bonds form to some extent, resulting in a downfield shift of the phenolic proton to 5.84 ppm. In an

unpolar solvent, such as benzene-*d*₆, the hydrogen bonding becomes more pronounced, shifting the phenolic proton down to 11 ppm.

In CHCl_3 , the target phenols **1a,b** showed broad, down-field peaks for the phenolic protons at 10 ppm for **1a** and at 10.7 ppm for **1b**. This strongly indicates a high degree of *intramolecular* hydrogen bonding, as suggested by our calculations. Further studies of the properties of these interesting compounds are underway and will be reported separately.

To summarize, novel 2-substituted imidazoles could be synthesized by a palladium-catalyzed cross-coupling between benzylzinc reagents and 2-iodo-imidazoles. The coupling is a promising alternative to existing routes towards these heterocycles, and is insensitive to steric hindrance in the benzylic moiety whereas the substituent in the 1-position of the imidazole seems important.

Preparation of zinc reagents. Synthesis of 2-methoxybenzylzinc bromide (7a): Zn-dust (1.48 g, 22.7 mmol) was suspended in anhydrous THF (5 mL) in a dried three-neck flask and heated to 60 °C under inert atmosphere. 1,2-Dibromoethane (0.08 mL, 0.93 mmol) was added and the mixture was stirred for 15 min at 60 °C. The flask was cooled to rt and chlorotrimethylsilane (0.1 mL, 0.78 mmol) was added. The mixture was stirred for 30 min at rt, then a solution of 2-methoxybenzyl bromide (3.8 g, 18.9 mmol) in THF (20 mL) was added over 2 h and the flask was sealed and stored in the fridge over night to let the Zn-dust settle. The concentration of the Zn reagent was not determined; 100% conversion was assumed. The solution could be stored in the fridge and the supernatant liquid was subsequently used in the Pd-coupling.

General Pd-coupling procedure. Synthesis of 2-(2-methoxy-benzyl)-1-methylimidazole (9a): $\text{Pd}(\text{OAc})_2$ (15 mg, 0.068 mmol) and tri(*o*-tolyl)phosphine (41 mg, 0.13 mmol) were dissolved in anhydrous THF (1 mL) under inert atmosphere. The mixture was stirred for 5 min before addition of imidazole **5a**¹⁴ (0.28 g, 1.35 mmol). The reaction mixture was cooled to 0 °C and the solution of **7a** (9 mL, 5 equiv.) was added over 5 min, and then the reaction was stirred at rt over night. Water and chloroform were added and the pH of the aqueous layer was adjusted to ~14 with NaOH (aq). The aqueous layer was extracted with 3* CHCl_3 , the combined organic layers were washed with 1M NaOH, dried with MgSO_4 and the solvent was evaporated. Column chromatography of the residue with EtOAc:MeOH 14:1 afforded **9a** as a pale yellow oil (0.16 g, 59%). ¹H NMR (400 MHz, CDCl_3): δ 7.20 (dt, 1H, *J*=7.7; 1.6 Hz), 6.98-6.96 (m, 2H), 6.89-6.85 (m, 2H), 6.80 (s, 1H), 4.07 (s, 2H), 3.85 (s, 3H), 3.48 (s, 3H); ¹³C NMR (100 MHz, CDCl_3): δ 156.5, 146.6, 129.9, 127.7, 127.2, 125.7, 120.6, 120.6, 110.2, 55.3, 32.6, 26.7; IR (in CH_2Cl_2): 3054, 2987, 1494, 1421 cm^{-1} .

2-(1-Methyl-imidazol-2-ylmethyl)-phenol (1a): mp 171.5-172.5 °C; ¹H NMR (300 MHz, CDCl_3): δ 11.0-9.0

(br s, 1H), 7.15 (t, 1H, $J=7.8$ Hz), 7.07 (d, 1H, $J=7.2$ Hz), 6.97 (d, 1H, $J=7.8$ Hz), 6.90 (s, 1H), 6.80 (t, 1H, $J=7.2$ Hz), 6.73 (s, 1H), 4.00 (s, 2H), 3.67 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3): δ 157.3, 147.8, 129.9, 129.1, 126.3, 124.0, 120.7, 120.1, 119.4, 33.2, 30.2; IR (KBr tablet): 2940, 1594, 1501, 1455 cm^{-1} .

2,4-Di-tert-butyl-6-(1-methyl-imidazole-2-ylmethyl)-phenol (1b): mp 158-161 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3): δ 10.7 (br s, 1H), 7.24 (d, 1H, $J=2.4$ Hz), 6.97 (d, 1H, $J=2.4$ Hz), 6.88 (d, 1H, $J=1.4$ Hz), 6.71 (d, 1H, $J=1.4$ Hz), 3.98 (s, 2H), 3.70 (s, 3H), 1.43 (s, 9H), 1.29 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3): δ 153.7, 148.2, 141.6, 138.8, 126.2, 124.5, 124.1, 123.5, 120.6, 35.4, 34.4, 33.2, 31.9, 30.8, 30.0; IR (KBr tablet): 3118, 1677, 1584, 1526 cm^{-1} .

2-(3,5-Di-tert-butyl-2-methoxy-benzyl)-1-methylimidazole (9b): mp 99-100 $^{\circ}\text{C}$; ^1H NMR (300 MHz, CDCl_3): δ 7.18 (d, 1H, $J=2.7$ Hz), 6.96 (d, 1H, $J=1.5$ Hz), 6.78 (m, 2H), 4.15 (s, 2H), 3.79 (s, 3H), 3.28 (s, 3H), 1.40 (s, 9H), 1.19 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3): δ 155.3, 147.5, 146.1, 141.7, 130.2, 127.4, 125.2, 122.7, 121.0, 61.9, 35.5, 34.6, 32.8, 31.6, 31.3, 28.4; IR (KBr tablet): 3102, 2961, 1500, 1477 cm^{-1} .

Acknowledgment

The Swedish National Energy Agency and the Swedish Foundation for Strategic Research (Selchem program) are gratefully acknowledged for funding this study.

References

- (1) Pesavento, R. P.; Van Der Donk, W. A. *Adv. Protein Chem.* **2001**, 58, 317 and references cited therein.
- (2) Stryer, L. *Biochemistry*, 5th ed.; W. H. Freeman and Company, New York, 1995.
- (3) Hays, A.-M. A.; Vassiliev, I. R.; Golbeck, J. H.; Debus, R. J. *Biochemistry* **1999**, 38, 11851.
- (4) The calculations will be reported separately together with further analyses of the target compounds.
- (5) Frutos, R. P.; Gallou, I.; Reeves, D.; Xu, Y.; Krishnamurthy, D.; Senanayake, C. H. *Tetrahedron Lett.* **2005**, 46, 8369 and references cited therein.
- (6) Abbotto, A.; Bradamante, S.; Pagani, G. A. *J. Org. Chem.* **1996**, 61, 1761.
- (7) Abarca-Gonzalez, B.; Jones, R. A.; Medio-Simon, M.; Quilez-Pardo, J.; Sepulveda-Arques, J.; Zaballos-Garcia, E. *Synth. Commun.* **1990**, 20, 321.
- (8) Moreno-Manas, M.; Bassa, J.; Llado, N.; Pleixats, R. *J. Het. Chem.* **1990**, 27, 673.
- (9) Murahashi, S.-I.; *J. Organomet. Chem.* **2002**, 653, 27.
- (10) Negishi, E.; Zeng, X.; Tan, Z.; Qian, M.; Hu, Q.; Huang, Z. In *Metal-Catalyzed Cross-Coupling Reactions*, 2nd ed., Vol. 2; De Meijere, A.; Diederich, F., Eds; Wiley-VCH Weinheim, 2004, 815-889.
- (11) Knochel, P.; Singer, R. D. *Chem. Rev.* **1993**, 93, 2117.
- (12) Angioletti, M. E.; Casalnuovo, A. L.; Selby, T. P. *Synlett* **2000**, 6, 905.
- (13) This reagent is also commercially available from Aldrich.
- (14) Prasad, A. S. B.; Stevenson, T. M.; Citineni, J. R.; Nyzam, V.; Knochel, P. *Tetrahedron* **1997**, 53, 7237.
- (15) Reiter, L. A.; Mitchell, P. G.; Martinelli, G. J.; Lopresti-Morrow, L. L.; Yocum, S. A.; Eskra, J. D. *Bioorg. Med. Chem. Lett.* **2003**, 13, 2331.
- (16) Compound **9c** was not deprotected due to lack of material.

Short title: Cross-Coupling of Imidazoles with Benzylzinc reagents